



Figure 3.2-32. Historical Snow Crab Catch (in 1,000 mt) in the Bering Sea. Some catches are given per season (spanning two calendar years), whereas others are per calendar year.




Source: ADF&G 2000b

Figure 3.2-33. Historical Korean Hair Crab Catch (in 1,000 mt) in the Bering Sea. Some catches are given per season (spanning two calendar years), whereas others are per calendar year.




Source: ADF&G 2000b

Figure 3.2-34. Historical Annual Dungeness Crab Catch (in 1,000 mt) in the PWS, Cook Inlet, Kodiak, and Alaska Peninsula Districts.




Source: ADF&G 2000b

Figure 3.2-35. Management Areas Involving Prohibited Species and Walrus in the Bering Sea




Source: NMFS 2001a

Figure 3.2-36. Historical Trend of Pacific Herring Catch in the Bering Sea and the Gulf of Alaska, 1990-1998.




Source: ADF&G Fish Ticket Landing Receipt Database, 1999

Figure 3.2-37. Final Critical Habitat for Spectacled and Steller's Eider




Source: NMFS 2001b

Figure 3.2-38. Distribution of Horny Corals (kilograms per hectare [kg/ha])




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-39. Distribution of Sponges (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-40. Distribution of Bryozoans (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-41. Distribution of Hydroids (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-42. Distribution of Sea Raspberries (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-43. Distribution of Sea Pen (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-44. Distribution of Anemones (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-45. Distribution of Sea Onion (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-46. Distribution of Sea Peach (kg/ha)



Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.3-1. Biomass Trends in Bering Sea Trophic Guilds, 1979 to 1998




Source: NMFS 2001b

Figure 3.3-2. Trophic Interactions of Key Eastern Bering Sea Groundfish




Source: NMFS 2001a

Figure 3.3-3. Trophic Relationships of the Groundfishes in the Aleutian Islands




Source: NMFS 2001a

Figure 3.3-4. Trophic Relationships of the Groundfishes in the Gulf of Alaska




Source: NMFS 2001a

Figure 3.3-5. Trend in Index of Species Composition Based on Ordination of Species Abundance Data from Five Triennial Surveys on Gulf of Alaska Shelf and Slope with Approximate 95 Percent Confidence Interval



Source: NMFS 2001a


Figure 3.4-1. Pelagic Trawl



Source: Jennings et al. 2001


Permission to use figure requested by Craig Rose (pending)

Figure 3.4-2. Bottom Trawl




Source: K. Richards, Tetra Tech FW, Bothell, WA

Figure 3.4-3. Set Longline Gear




Source: K. Richards, Tetra Tech FW, Bothell, WA

Figure 3.4-4. Vertical Longline




Source: K. Richards, Tetra Tech FW, Bothell, WA

Figure 3.4-5. Jig Machine



Source: K. Richards, Tetra Tech FW, Bothell, WA


Figure 3.4-6. Gulf of Alaska: 1990-2002 Observed Pollock Pelagic Trawl Effort (Sets/25 km²)



Note: (Sets/25 km²) = Summed for 1990 to 2002

Source: Cathy Coon, NPFMC Staff, North Pacific Observer Program Data


Figure 3.4-7. Gulf of Alaska: 1990-2002 Observed Pollock Non-Pelagic (Bottom) Trawl Effort (Sets/25 km²)



Note: (Sets/25 km²) = Summed for 1990 to 2002

Source: Cathy Coon, NPFMC Staff, North Pacific Observer Program Data


Figure 3.4-8. Gulf of Alaska: 1990-2002 Observed Pacific Cod Non-Pelagic (Bottom) Trawl Effort (Sets/25 km²)



Note: (Sets/25 km²) = Summed for 1990 to 2002

Source: Cathy Coon, NPFMC Staff, North Pacific Observer Program Data


Figure 3.4-9. Gulf of Alaska: 1990-2002 Observed Deepwater Flatfish Non-Pelagic Trawl Effort (Sets/25 km²)



Note: (Sets/25 km²) = Summed for 1990 to 2002

Source: Cathy Coon, NPFMC Staff, North Pacific Observer Program Data


Figure 3.4-10. Gulf of Alaska: 1990-2002 Observed Shallow Water Flatfish Non-Pelagic Trawl Effort (Sets/25 km²)



Note: (Sets/25 km²) = Summed for 1990 to 2002

Source: Cathy Coon, NPFMC Staff, North Pacific Observer Program Data

Figure 3.4-11. Gulf of Alaska: 1990-2002 Observed Slope Rockfish Pelagic Trawl Effort (Sets/25 km²)



Note: (Sets/25 km²) = Summed for 1990 to 2002

Source: Cathy Coon, NPFMC Staff, North Pacific Observer Program Data